

INTEGRATED WEED MANAGEMENT IN DIFFERENT VARIETIES OF ONION (*Allium cepa* L.)

Abdul Ghaffoor¹

ABSTRACT

Three onion varieties viz. Naurang, Dark Red and Swat-1, two chemical treatments (2,4-D @ 2.5 and pendimethaline @ 3.0 L ha⁻¹) and two manual weeding treatments (Single hoeing 25 days and Double hoeing 25 & 50 days after transplanting) alongwith a weedy check were compared for weed control and their effect on the yield and yield components of onion. Varieties were assigned to the main-plots while weed control treatments were kept into subplots. Each sub-plot size was of 1.5 x 1.5 m² having the distance of 30 cm and 10 cm between rows and plants, respectively. Swat-1 showed the best results in all the parameters including number of leaves plant⁻¹, length of leaves, number of bulbs plots⁻¹, bulb diameter, weight of bulbs and bulb yield (t ha⁻¹). Weedicides significantly controlled the weeds. However, pendimethaline @ 3 L ha⁻¹ treated plots gave the highest yield (40.28 t ha⁻¹) as well as weight of bulbs (127.9 g). Our findings reveal that Swat-1 variety is suitable for the area under the climatic conditions of Dera Ismail Khan, whereas pendimethaline proved to be the best weedicide in controlling weeds in onion. Therefore, it is recommended to have a combination of Swat-1 and pendimethaline for the higher yield of onion for Dera Ismail Khan area.

Key words: Varieties, weed control methods, 2,4-D, pendimethaline, onion.

INTRODUCTION

Pakistan is an agricultural country with its economy being greatly dependent upon agriculture. Although, the major contributors to the national economy are the major crops such as cotton, rice and wheat yet the production of minor crops like onion (*Allium cepa* L.) also has a pronounced impact on the economy of the country. Onion belongs to the lily family or Alliaceae and is a condiment crop and consumed as a fresh in salads as well as used in dishes as a spice. Besides providing nutrition, it imparts acceptable flavor to our dishes. The yield of onion in our country is very low as compared to other agricultural countries. One of the most important reasons for the poor yield of onions is the excessive weed competition. The most comprehensive definition of weed is the plant whose virtues have not been discovered so far. Weed control is unavoidable for successful production of the onion crop and production losses increased with weed infestation. Weed control constitutes one of the principal costs of production in onions (Nyland, et al., 1958). Orkwoj et al., (1983) reported that Chlorthal dimethyl @ 12 kg ha⁻¹, flurodifen @ 1.5 and 3 kg ha⁻¹ and oxadiazon @ 1 and 2 kg ha⁻¹ all applied prior to transplanting gave excellent weed control for at least 12 weeks and resulted in onion yield comparable with hoe weeding. Herbicides applied after transplanting were not so satisfactory. Orkwoj (1983) reported that weed infestation significantly reduced crop

1 Department of Horticulture, Faculty of Agriculture, Gomal University, Dera Ismail.Khan, NWFP
– Pakistan E-mail:waseem_khan49@hotmail.com.

vigor, leaf production and bulb diameter and consequently bulb yield in onions. Patel *et al.*, (1986) reported that Fluchloralin @ 1.35 kg ha⁻¹ plus one hand weeding gave the highest additional income compared with no weeding. The cost benefit ratio was highest with 0.5 kg oxidiazon ha⁻¹. Rajendra *et al.*, (1986) concluded that nitrofen @ 1 g ha⁻¹ applied pre-planting gave good weed control and highest yields of 282.41 g ha⁻¹ compared with 193.21 g ha⁻¹ in the control. Nitrofen treatment was at par with 2-hand weeding and it was more economical. Plots hand weeded 3 and 4 times yielded 302.79 and 309.26 g ha⁻¹, respectively. The critical weed competition was up to 40 days after transplanting. Porwal and Singh (1993) observed that the pre-emergence application of oxadiazon (1.0 kg ha⁻¹) followed by manual weeding 45 days after transplanting showed the maximum weed-control efficiency of 77.0% compared with 60.5% with oxadiazon alone, and also increased the bulb yield by 81.4-254.9% over weedy check. Warade *et al.*, (1995) obtained highest yields of 11.83 t ha⁻¹ and largest bulbs were achieved with 0.5 kg fluchloralin, followed by 0.75 kg fluchloralin. The highest return of Rs.35499/- and best benefit cost ratio (25.46) were also achieved with 0.5 kg fluchloralin. Verma and Singh (1997) reported higher plant height, leaves plant⁻¹, fresh and dry weight of plant (g plant⁻¹), bulb diameter, marketable bulb yield and poor percentage of bolting recorded under weed free plot, followed by pendimethalin at 1.5 kg ha⁻¹ and at higher fertility.

MATERIALS AND METHODS

The research project on Integrated Weed Management in Different Varieties of Onion (*Allium cepa* L.) was carried out at the Horticultural Research Area, Faculty of Agriculture, Gomal University, Dera Ismail Khan, during the growing season 2001-2002. The experiment was laid out in split plot design with three replications, allotting the varieties (Naurang, Dark Red and Swat-I) into main plots while weeding methods into sub plots. The sub-plots consisted of herbicide treatments viz. Weedy Check (Control), Single Hoeing (25 Days after Sowing), Double Hoeing (25 and 50 Days after Sowing), 2, 4-D @ 2.5 L ha⁻¹ and pendimethaline @ 3.0 L ha⁻¹. Each sub-plot size was of 1.5 x 1.5 m² having the distance of 30 cm and 10 cm between rows and plants, respectively. 2,4-D was applied in post emergence of weeds whereas pendimethaline was sprayed immediately after transplanting. First hand weeding was applied 25 days after transplanting while the second hand weeding was applied with 25 days interval. Data were recorded on bolting plot⁻¹ (%), weed density m⁻², fresh weeds biomass (g plot⁻¹), dry weeds biomass (g plot⁻¹), number of leaves plant⁻¹, number of bulbs plot⁻¹, diameter of bulbs (cm), weight of bulb (g) and bulb yield (t ha⁻¹). Data were subjected to ANOVA and the significant means were separated as outlined by Steel and Torrie, 1984.

RESULTS AND DISCUSSION

Weed density 25 days after transplanting (m⁻²)

Table-1 shows weed density was not affected significantly in case of three onion varieties. However, lesser number of weeds (191.2 m⁻²) were counted in the plots where Dark Red variety was sown which were very closely followed by Swat-I and Naurang with 191.3 and 191.9 weeds m⁻², respectively. Whereas, different weeding methods gave significant results for the weed density. Lower weed count was recorded in herbicides treated plots where 120.9 and 125.3 weeds m⁻² were recorded in pendimethaline and 2,4-D treated plots, respectively. Statistically both the treatments were at par with each other. Maximum number of weeds per m² were counted as 296.7 from weedy check plots. This might be due to the application of herbicides.

Fresh weed biomass (g)

The data showed that different onion varieties differed non-significantly (Table-1). However, maximum weight of weeds was measured from plots where Naurang was sown with 321.90 g weeds biomass plot⁻¹, whereas variety Dark Red gave 268.5 g weeds biomass plot⁻¹ as the minimum. Different weed control methods showed a significant variation. Control plots having no weed control produced the maximum weed weight i.e. 461g followed by double and single hoeing treatments with 442.40 & 350.20 g respectively. All these three treatments were statistically at par with each other. Pendimethaline showed the minimum weight of the weeds (66.52 g). The results are in accordance with the findings of Verma and Singh (1997) who stated that pendimethaline controlled the weeds biomass.

Dry weeds biomass (g)

Dry weed biomass was significantly affected by the three onion varieties. Minimum dry weed biomass (22.43 g) was recorded from Dark Red variety. The plots of Naurang variety produced the maximum dry weed biomass i.e. 48.30 g. Different weed control practices showed the significant variations for the said parameter. The minimum dry weed biomass (23.98 g) was observed in pendimethaline treated plots. The plots with Single Hoeing gave the maximum dry weed biomass i.e. 42.81 g followed by control plots (38.97 g). Statistically the results of Control, Single Hoeing and Double Hoeing were comparable with one another (Table-1). Verma and Singh (1997) also reported that chemicals controlled the fresh as well as dry weed biomass.

Number of leaves per plant

The results on leaves per plant revealed the significant differences for both factors. Maximum leaves plant⁻¹ (10.32) were obtained in Swat-I variety while Dark Red gave the lowest leaves plant⁻¹ i.e. 8.56. This might be due to the varietal characteristics of the concerned cultivar. Different weed control methods also showed significant variations. Maximum number of leaves plant⁻¹ was recorded as 11.57 from pendimethaline treated plots. While the lowest leaves plant⁻¹ producing treatment was Control (8.04) where no weeding was done (Table-1). Disappearance of weeds resulted in the better fertilizer use efficiency by the plant and in this way the plants accumulated more food and increased the number of leaves compared to the treatments where hoeing was given.

Leaf length (cm)

Similar trend was observed as it was recorded for the number of leaves. Both the factors and their interaction showed significant differences. Maximum length of leaves (51.41 cm) was obtained from Swat-I while Dark Red variety produced the shortest leaves of 39.01 cm length. The plants of various varieties utilized the plant food nutrients easily without any competition which resulted in better length of leaves. Whereas, pendimethaline treated plots gave the longest leaves of onion i.e. 49.42 cm against the control plots which produced minimum leaf length (41.12 cm). Interaction of both factors also revealed significant variations. Swat-I with the treatment of Single Hoeing and pendimethaline gave the longest leaf length viz. 53.37 cm and 53.27 cm, also non-significant with each other (Table-1). Verma and Singh (1997) also reported the same findings during their field trial.

Number of bulbs per plot

Onion bulbs are the economical and most important commodity. Both factors showed the significant behavior with regards to number of bulbs plot⁻¹. However, Swat-I showed the best results by producing 67.55 bulbs plot⁻¹, while Dark Red was the minimum bulb producing variety giving 64.47 bulbs plot⁻¹. As the plots treated with pendimethaline remained free of weeds, they produced more number of bulbs (69.78) without any difficulty and deficiency of nutrients. Statistically the results of pendimethaline (69.78 bulbs plot⁻¹), Double Hoeing (67.89 bulbs plot⁻¹) and Single Hoeing (66.78 bulbs plot⁻¹) were non-significant among one another, and remained at par with each other. Minimum bulbs plot⁻¹ (62.00) was recorded in the Control plots (Table-2). These findings go in accordance with those of Porwal and Singh (1993) who stated that weeds control treatments reduced the weeds density and enhanced the bulb yield.

Diameter of bulbs (cm)

Table-2 depicted that different varieties of onion significantly affected the diameter of onion bulbs. Swat-I was the maximum bulb diameter producing variety giving 6.49 cm wide bulbs. The minimum diameter was noted in Dark Red variety i.e. 6.18 cm. Various weed control measures also significantly affected the bulb diameter. The plots treated with pendimethaline and hand weeding (Hoeing) increased the bulb diameter of onion. Pendimethaline treated plots produced 6.63 cm wide bulbs showing the maximum bulb diameter. Minimum bulb size was noted as 5.96 cm from weedy check plots where the weed control practices were not applied. Pendimethaline and Single Hoeing practice gave at par findings and also remained non-significant with each other. Chemically treated plots remained non-competitive with weeds throughout the growing season which resulted more wide onion bulbs.

Weight of bulbs (g)

The three onion cultivars differed significantly for the weight of bulbs (Table-2). Swat-I produced the maximum weighing bulbs (122.30 g) whereas the lowest weight of bulbs was recorded as 105.30 g from the Dark Red variety. Different weed control practices also affected significantly the weight of onion bulbs. Plots treated with pendimethaline showed maximum bulb weight i.e. 127.90 g as against the minimum of 105.10 g (Control) which was very closely followed by 2,4-D treated plots with 106.20 g bulb weight. The results coincide with the findings of Dunan *et al.*, (1996).

Bulb yield (t ha⁻¹)

The three varieties significantly affected the onion bulb yield. Swat-I was the highest yield producing variety with 36.76 t ha⁻¹ yield. The lowest yield was recorded as 30.21 t ha⁻¹ from Dark Red variety. Pre-emergence application of pendimethaline boosted up the onion bulb yield. Whereas, pendimethaline treated plot produced maximum yield of 40.08 t ha⁻¹ while the minimum of 29.08 t ha⁻¹ from the weedy check plots (Table-2). These results are also supported by the findings of Warade *et al.*, (1995) and Shimi and Moghadam (1996) who stated that herbicides application significantly increased the bulb yield of onion.

Table-1. Weeds density (25 DAS), fresh weed biomass (g), dry weed biomass (g), number of leaves plant⁻¹ and leaf length (cm) of three onion varieties as affected by various weed control methods

Weeding Methods	Varieties			Weed Management Mean
	Naurang	Dark Red	Swat-I	
Weeds Density 25 Days After Sowing				
Control (No Weeding)	303.33 NS	298.33	288.33	296.67 a
Single Hoeing (25 DAS)	201.67	211.67	215.00	209.44 b
Double Hoeing (25,50 DAS)	209.33	195.00	210.33	204.89 b
2,4 - D (1 L ha ⁻¹)	125.00	127.67	123.33	125.33 c
Pendimethaline (1.25 L ha ⁻¹)	120.00	123.33	119.33	120.89 c
Varietal Means	191.87 NS	191.20	191.27	
Fresh Weed Biomass (g)				
Control (No Weeding)	575.90 NS	522.80	284.37	461.00 a
Single Hoeing (25 DAT)	441.90	298.00	310.67	350.20 a
Double Hoeing (25,50 DAT)	365.90	355.17	606.20	442.40 a
2,4 - D (1 L ha ⁻¹)	147.23	119.90	138.10	135.10 b
Pendimethaline (1.25 L ha ⁻¹)	078.61	046.67	074.28	066.52 b
Varietal Mean	321.90 NS	268.50	282.70	
Dry Weed Biomass (g)				
Control (No Weeding)	25.40 NS	27.37	64.07	38.97 a
Single Hoeing (25 DAT)	35.93	28.70	63.80	42.81 a
Double Hoeing (25,50 DAT)	47.50	21.80	44.37	37.89 a
2,4 - D (1 L ha ⁻¹)	28.73	21.50	37.37	27.87 b
Pendimethaline (1.25 L ha ⁻¹)	23.27	12.70	35.90	23.98 c
Varietal Mean	32.18 b	22.43 b	48.30 a	
Number of Leaves per Plant				
Control (No Weeding)	08.23 NS	07.83	08.07	08.04 c
Single Hoeing (25 D DAT)	08.70	08.17	10.23	09.03 bc
Double Hoeing (25,50 DAT)	09.03	09.27	11.73	10.01 b
2,4 - D (1 L ha ⁻¹)	09.26	07.83	08.57	08.56 c
Pendimethaline (1.25 L ha ⁻¹)	12.00	09.70	13.00	11.57 a
Varietal Mean	09.45 b	08.56 c	10.32 a	
Leaf Length (cm)				
Control (No Weeding)	38.50 e	36.10 e	48.77 bcd	41.12 d
Single Hoeing (25 DAT)	49.33 bcd	36.17 e	51.73 ab	45.74 bc
Double Hoeing (25,50 DAT)	49.50 bcd	38.67 e	53.37 a	47.18 b
2,4 - D (1 L ha ⁻¹)	45.93 d	37.57 e	49.90 abc	44.47 c
Pendimethaline (1.25 L ha ⁻¹)	48.43 bcd	46.57 cd	53.27 a	49.42 a
Varietal Mean	46.34 b	39.01 c	51.41 a	

Any two mean(s) in their respective group not sharing a common letter(s) are significant at 5% probability.

NS = Non-significant

DAT= Days after transplanting

Table-2. Number of bulbs plot⁻¹, diameter of bulbs (cm), weight of bulbs (g) and yield (t ha⁻¹) of three onion varieties as affected by various weed control methods

Weeding Methods	Varieties			Weed Management Mean
	Naurang	Dark Red	Swat-I	
Number of bulbs plot⁻¹				
Control (No Weeding)	62.33 NS	60.00	63.67	62.00 c
Single Hoeing (25 DAT)	67.00	65.33	68.00	66.78 ab
Double Hoeing (25,50 DAT)	67.33	67.33	69.00	67.89 a
2,4 - D (1 L ha ⁻¹)	64.00	63.67	64.67	64.11 bc
Pendimethaline (1.25 L ha ⁻¹)	71.00	66.00	72.33	69.78 a
Varietal Mean	66.33 ab	64.47 b	67.53 a	
Diameter of the Bulbs (cm)				
Control (No Weeding)	5.93 NS	5.73	6.20	5.96 c
Single Hoeing (25 DAT)	6.37	6.43	6.57	6.46 ab
Double Hoeing (25,50 DAT)	6.40	6.30	6.50	6.40 b
2,4 - D (1 L ha ⁻¹)	6.03	5.90	6.43	6.12 c
Pendimethaline (1.25 L ha ⁻¹)	6.63	6.53	6.73	6.63 a
Varietal Mean	6.27 ab	6.18 b	6.49 a	
Weight of Bulbs (g)				
Control (No Weeding)	105.30 NS	94.33	115.70	105.10 c
Single Hoeing (25 DAT)	115.70	101.30	123.00	113.30 b
Double Hoeing (25,50 DAT)	121.30	109.70	124.70	118.60 b
2,4 - D (1 L ha ⁻¹)	116.30	93.33	119.00	106.20 c
Pendimethaline (1.25 L ha ⁻¹)	128.00	126.30	129.30	127.90 a
Varietal Mean	115.30 b	105.00 c	122.30 a	
Bulb Yield (t ha⁻¹)				
Control (No Weeding)	29.43 NS	25.17	32.64	29.08 c
Single Hoeing (25 DAT)	34.43	29.46	37.18	33.69 b
Double Hoeing (25,50 DAT)	36.31	32.75	38.24	35.77 b
2,4 - D (1 L ha ⁻¹)	30.28	24.73	34.19	29.73 c
Pendimethaline (1.25 L ha ⁻¹)	40.39	38.92	41.54	40.28 a
Varietal Mean	34.17 a	30.21 b	36.76 a	

Any two mean(s) in their respective group not sharing a common letter(s) are significant at 5% probability.

NS = Non-Significant

DAT = Days After Transplanting

REFERENCES CITED

- Dunan, C.M., P. Westra, F. Moore and P. Chapman. 1996. Modeling the effect of duration of weed competition, weed density and weed competitiveness on seeded and irrigated onion. *Weed Res.* 36(3): 259-269.
- Nyland, R.E., D.C. Nelson, and D.H. Dinkel. 1958. Comparative costs of weeding onions by hand or with monuron, CIPC and CDAA. *Weeds* 6:304-309.
- Orkwor, G.C. 1983. Effect of herbicides on the agronomic characters that control bulb yield. *Hortic. Absts.* 53(5): 3254; 1983.
- Orkwor, G.C., M.K. Moolani, A.M. Choudhary. 1983. A study of weed control in irrigated onions in Northern Nigeria. *Hortic. Absts.* 53(5): 3253; 1983.
- Patel, C.L., Z.G. Patel and R.B. Patel. 1986. Efficiency and economics of some new herbicides for weed control in onions. *Gujarat Agric. Univ. Res. J. India.* 12(1):14-17. *Hortic. Absts.* 58(1-7): 864;1986.
- Porwal, M.K. and M.M. Singh. 1993. Effect of N and weed management on onion. *Indian J. Agron.* 38(1): 74-77.
- Rajendara, S., A.P. Sinha and S.P. Singh. 1986. Comparative efficacy of herbicides and hand weeding for the control of weeds in onions. *Haryana J. Hort. Sci.* 15(1-2): 76-77.
- Shimi, P. and R.N. Moghadam. 1996. Investigation on the efficacy of 2 herbicides in onion fields. *Seed and Plant* 12(3): 34-37.
- Steel, R.G.D. and J. H. Torrie. 1984. Principles and procedures of statistics, a biological approach. Mc Graw Hill Book Co. Inc. New York.
- Verma, S.K. and T. Singh. 1997. Effect of weed control measures and fertility on growth and productivity of rainy season onion. *Indian J. Agron.* 42(3): 540-543.
- Warade, S.D., S.B. Desale and K.G. Shinde. 1995. Effects of weedicides on weed intensity and yield of Kharif Onion cv. Phule Safed. *J. Maharashtra Agric. Univ.* 20(3): 462-463.